

# The of Ciplukan, Betel, Moringa and Patchouli Waste Extract as Phytogetic Feed Additive and Their Effect on Broiler Performance

M Daud<sup>1\*</sup>, M A Yaman<sup>1</sup>, Y Usman<sup>1</sup>, C R Maulida<sup>1</sup>

<sup>1</sup>Animal Husbandry, Faculty of Agriculture, Syiah Kuala University, Banda Aceh, 23111 Indonesia

\*Corresponding author. Email: [daewood@unsyiah.ac.id](mailto:daewood@unsyiah.ac.id)

## ABSTRACT

The aim of this study was to examine the potential of ~~ciplukan (*Physalis angulata L.*), betel (*Piper betle L.*), moringa (*Moringa oliefera L.*) and patchouli (*Pogostemon patchouli Pellet*)~~ waste as a phytogetic feed additive to replace Antibiotic Growth Promoter (AGP) and their effect on broiler performance. The study was conducted experimentally, using a completely randomized design (CRD) consisting of 5 treatments ~~and with~~ 4 replications. The treatments ~~consisted of used were~~: R0 (control), R1 (0.5% ciplukan extract), R2 (0.5% betel extract), R3 (0.5% ~~m~~Moringa extract) and R4 (0.5% patchouli waste extract). ~~The variables observed were~~: aAntioxidant content, ration consumption, final body weight, ration conversion and broiler mortality ~~were observed~~. The ~~obtained data obtained~~ were analyzed by Analysis of Variance (ANOVA) and continued with Duncan's multiple distance test. The results showed that the highest antioxidant content was found in ciplukan extract ~~, namely (-75.69%) followed by the~~ patchouli extract (73.53%), ~~Moringa moringa~~ extract (73.41%), and then the lowest ~~was~~ betel extract (61.29%). ~~Giving~~ Addition of ciplukan, betel, moringa and patchouli waste extracts with a percentage of 0.5% in drinking water did not significantly affect ration consumption, final body weight, ration conversion and broiler mortality. It was concluded that the highest antioxidant content was obtained in ciplukan extract and the addition of 0.5% ciplukan, betel, moringa and patchouli waste extract in drinking water could ~~have not a negatif~~ ~~positive~~ effect on the appearance of broiler.

## Keywords:

## 1. INTRODUCTION

One of the obstacles that are often faced in poultry farming is the problem of decreasing growth performance and increasing disease ~~attacks~~infections. Efforts that can be made to increase productivity and prevent disease ~~attaeks~~infections in poultry are by increasing non-specific immune responses through the provision of feed additives. Feed additives that are very commonly used, are antibiotics or better known as Antibiotic Growth Promoter (AGP), which serves to help fight pathogenic bacteria and consequently can increase livestock production. However, the use of antibiotics today has received special attention from the government and consumers. Due to the administration of certain antibiotics, livestock are resistant to Enrofloxacin which functions to eradicate *Escherichia*

*coli* ~~bacteria~~. WHO has banned the use of AGP in livestock because it can be bad for human health.

In connection with the prohibition on the use of ~~antibiotic growth promoters~~ (AGP) in poultry feed, it is necessary to find alternative livestock growth promoters from other sources, in addition to supporting food safety, also having the ability to act as antibiotics in increasing livestock productivity and health. One alternative growth promoter that can be used today in poultry rations is phytogetic as a feed additive. Phytogetic feed additives (phytobiotics or botanicals) are generally defined as additional compounds, which are the result of plant secondary metabolites. They either contain nutritional, non-nutritious, or anti-nutritional compounds that are included in the ration to increase livestock productivity by improving feed properties and the qualities of livestock production, as

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Commented [D1]: Please add the scientific name

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: 10 pt

Commented [D2]: This is doubt statement since in the previous sentence the author mention that those by-products has no effect on broiler.

Formatted: Font: Italic

well as increasing production performance and digestive tract health by controlling pathogenic bacteria [1], [2].

Feed additives are generally considered as products applied to livestock, which aims at improving their health status and production performance. In contrast, they are generally applied for prophylactic and therapeutic purposes in veterinary medicine for health problems that have been diagnosed for a limited or a waiting period basis. However, phytochemicals and their application aspects are a relatively new group in feed additives. One of the herbal plants that are used as a source of phytochemical additive feeds is ciplukan (*Physalis angulata L.*), betel (*Piper betle L.*), moringa (*Moringa oleifera L.*), and patchouli (*Pogostemon patchouli Pellet.*) oil wastes. They have a height of 0.1-1 m, grow wildly and evenly in lowland areas to an altitude of 1.550 above sea level [3], and have a high nutritional value, which is rich in vitamins, minerals, and antioxidants [4]. Likewise, patchouli is an essential oil-producing plant that make an important contribution in the world of flavour and fragrance, especially for the perfume and aromatherapy industry. Meanwhile, this plant is used as a drug that functions as an anti-inflammatory, anti-depressant, divertic, antifungal, and antibacterial in the pharmaceutical industry [5].

However, there are not much information and reports on the use of ciplukan, betel, moringa and patchouli oil refined waste, which have the potential to be used as a source of feed additives for livestock. With this in mind, it is necessary to investigate the potential of these plants, such that it is used as an alternative source of phytochemical feed additives, in an effort to providing a replacement for Antibiotic Growth Promoter (AGP) in livestock. Therefore, this research aims of this study was to examine the potential of ciplukan, betel, moringa and patchouli oil waste as a phytochemical feed additive to replace Antibiotic Growth Promoter (AGP) and their effect on broiler performance.

## 2. MATERIALS AND METHOD

### 2.1 Material/sample Handling

Leave samples of ciplukan, betel, moringa plants and patchouli wastes was cleaned and dried by aerating for 7 days. Furthermore, these samples were mashed using a diskmill machine to produce mesh-shaped leaves and then stored for further testing.

### 2.2 Extraction of Ingredients

Meshed samples of Ciplukan, betel, moringa and patchouli wastes were extracted by the maceration method, which is used to attract the required compounds by immersing the material into

96% ethanol solvent. The use of this solvent is intended to extract chemical components and to determine the yield and antioxidant content of each sample.

Maceration extraction was carried out with a sample and solvent ratio of 1:4 for 48 hours at a 28°C. Subsequently, the extraction results were filtered with a vacuum filter paper. Afterward, the filtrate was evaporated using a rotary evaporator such that it gets a concentrated extract and is then stored at 0°C for the next test process.

The crude extract was weighed to determine the yield based on the extraction method, and the type of solvent using the formula:

$$\text{Yield} = \frac{\text{dry extract weight (g)}}{\text{initial sample weight (g)}} \times 100\%$$

### 2.3 Broiler Treatment

This study used 100 broiler chickens that were reared up to 4 weeks of age in 20 postal/litter cages with the size of each cage unit being 1 x 1 meter and covered with a pad of rice husk as thick as ± 10 cm. Each cage unit consisted of 5 broilers. The study was conducted using an experimental method, using a completely randomized design (CRD) with 5 ration treatments and 4 replications in quadruplicate.

The treatments consisted of: R0 (control), R1 (0.5% ciplukan extract), R2 (0.5% betel extract), R3 (0.5% moringa extract), R4 (0.5% patchouli waste extract). The observed variables were: antioxidant content, ration consumption, final body weight, ration conversion and broiler mortality. The obtained data were analyzed by Analysis of Variance (ANOVA) and continued with Duncan's multiple distance test.

## 3. RESULT AND DISCUSSION

### 3.1 The Antioxidant Content of Ciplukan, Betel, Moringa and Patchouli Waste Extract

Further test results showed that the antioxidant levels tended to be higher in the ciplukan extract with a percentage of 75.70%, followed by patchouli waste (patchouli leaves) of 73.53%, moringa extract of 73.41%, and betel extract of 61.29%, moringa extract of 73.41% and patchouli waste (patchouli leaves) of 73.53% (Table 1).

Table 1. Antioxidant test results.

Material/sample type	Antioksidan (%)
Ciplukan extract	75.70±0.66

Formatted: Indent: First line: 0,51 cm, Space After: 0 pt

Formatted: Indent: First line: 0 cm

Commented [D3]: Add scientific name

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Betel extract	61.29 ±9,94
Moringa extract	73.41 ±0,86
Patchouli waste extract	73.53±0.17

Source: Lab. of Food and Agricultural Product Analysis, Syiah Kuala University (2020).

Antioxidant is a substance that neutralizes or reduces the negative impact of free radical, which is a molecule containing the collection of an unpaired electron in an outer circle. Various researches have studied broadly the benefits of antioxidants to ward off free radicals. The results showed that antioxidants slow down processes caused by free radicals, such as the presence of tocopherol, ascorbate, flavonoids, and lycopene [6].

## 3.2. Broiler Performance

### 3.2.1 Ration Consumption

The consumption of broiler rations from each treatment is shown in Table 2. The results of statistical analysis showed that the consumption of broiler chicken rations ~~from all treatments there was had~~ no significant difference between treatments. ~~Relatively higher ration consumption was found in treatment T3 (moringa extract), but it was not significantly different when compared to other treatments (Table 2). However, the consumption of rations tended to be higher in the T3 treatment (moringa extract) when compared to other treatments.~~

~~Previous study Stated-stated~~ that the provision of ~~Moringa-moringa~~ leaves in rations up to a level of 24% did not have a significant effect on ration consumption, weight gain, and efficiency of ration use in chickens [7]. Furthermore, the results of research [8] also showed that giving betel leaf extract in drinking water up to 2% had no effect on ration consumption, daily body weight gain, feed conversion and final body weight of broiler chickens

### 3.2.2 Final Body Weight

The results showed that the administration of ciplukan leaf extract, betel leaf, moringa and patchouli leaf wastes did not significantly affect the final body weight of broiler chickens aged 4 weeks (Table 2).

However, the final body weight in T3 treatment with 0.5% ~~m~~Moringa leaf extract ~~tended~~ to be higher than other treatments.

Based on the research results of [9] that administration of 5% ~~m~~Moringa leaf extract through drinking water, was able to increase body weight gain and final body weight compared to controls because moringa leaves have high protein and carbohydrate content. In addition, the phytochemical compounds in

moringa leaves can inhibit the growth of harmful microorganisms in the digestive tract of chickens, so that the utilization of nutrients by broiler chickens can be optimal. As reported by [10], moringa leaves contain phytochemical compounds, such as flavonoids, saponins, tannins and several other phenolic compounds that have antimicrobial activity.

~~Several Research results [11]~~ also show that the administration of various herbs does not have a significant effect on livestock performance ~~[11]~~. ~~The similar result ame thing~~ was also reported ~~from the results of research conducted by by another study~~ [12] where broiler chickens ~~fed given~~ phytochemical feed additives (PFA) containing essential oils with different levels from 150-1500 mg/kg feed did not affect the weight gain of livestock at the age of 21, 35 or at the age of 42 days.

### 3.2.3 Feed Conversion Ratio

The results of statistical analysis showed that the cumulative conversion value of 4 weeks old broiler rations did not show a significant difference between treatments (Table 2). ~~The same thing~~ Similar results ~~was~~ as reported by [13], that the addition of boiled water of betel leaves into broiler drinking water did not have a significant effect on ration consumption, drinking water consumption and ration conversion, ~~and And also,~~ the research of [14] ~~reported that,~~ administration of 5% ~~Moringa-moringa~~ leaf extract and garlic leaves through drinking water ~~couldan~~ significantly increase the efficiency of the use of rations.

### 3.2.4 Mortality

The administration of ciplukan leaf extract, betel leaf, ~~Moringa-moringa~~ and patchouli leaf wastes did not show a significant difference in broiler mortality during the study. However, there were also deaths in the T3, T3, T4 and T0 treatments as shown in Table 2.

Based on the results of the necropsy test, the death of the chickens was caused by infection with the pathogenic Escherichia coli bacteria, which can cause abnormalities in chickens whose characteristics include: air sacs, severe heart bag inflammation and severe liver inflammation with a layer of fibrin that covers most or all of the surface of the liver with a grayish-white or yellowish color. Giving ciplukan extract (T1) there were no deaths in broiler chickens, this was because the active compounds contained in ciplukan extract could reduce mortality because these compounds had bacterial inhibitory power.

As the results of previous research [15] that ciplukan extract contains flavonoids and polyphenols, which are efficacious as antioxidants and antibacterials. The active

**Commented [D6]:** This mean to cite more than reference, but in the text only refer one reference. Please clarify it. References no 11 is not a review paper.

**Formatted:** Font: Times New Roman, 10 pt, Font color: Auto, Pattern: Clear

**Commented [D4]:** 'Tend' can be used if P-value was 0.05 < P < 0.10. Please consider to revise your discussion if the P-value is more than 0.10

**Commented [D5]:** Please see my previous comment

compounds contained in ciplukan ~~are include~~ saponins, flavonoids, polyphenols, chlorogenic acid, sugars and fisalin. Polyphenols and flavonoids are the most

abundant main compounds in ciplukan plants that can be used as antioxidants. Stated that flavonoids can inhibit the growth of bacteria [16].

**Table 2.** Performance of broiler chickens

Variable	Treatment				
	T0	T1	T2	T3	T4
Ration consumption (g/h)	2160.73±31.4	2113.10±30.5	2138.40±32.4	2360.98±38.7	2192.55±32.1
Final body weight (g/h)	1511.58±112.62	1590.45±37.72	1493.50±111.98	1620.12±43.97	1579.19±89.68
Feed conversion ratio	1.43±0.13	1.36±0.09	1.48±0.22	1.46±0.11	1.42±0.13
Mortality (%)	0.50±0.58	0.00±0.00	0.25±0.50	0.75±0.96	0.25±0.50

#### 4. CONCLUSION

It was concluded that the highest antioxidant content was obtained in ciplukan extract and the addition of 0.5% ciplukan, betel, moringa and patchouli waste extracts in drinking water could ~~not have~~ a ~~negative~~ ~~positive~~ effect on the appearance of broiler.

#### ACKNOWLEDGEMENTS

The authors are grateful to the research team, LPPM Syiah Kuala University, and all those that helped carry out the research.

#### REFERENCES

[1] Ganguly, S 2013 Phytogetic growth promoter as replacers for antibiotic growth promoter in poultry birds *Adv Pharmacoeipidem Drug Safety* **2** 2-3.

[2] Hashemi S R and Davoodi H 2011 *Vet Res Commun.* **35** 169-180.

[3] Monikawati A, Farida S, Putri L W, Ikhtisarsyah Y G and Meiyanto E 2011 *Indones J. Cancer Chemoprev* **2** 227-232.

[4] Wei J, Hu X and Yang X 2012 Indentification of single-copy orthologous genes between *Physalis* and *Solanum lycopersicum* and analysis of genetic diversity in *Physalis* using molocular markers. *Plos One.* 7(11), pp. e50164.

[5] Blank A F, Tricia C P S, Priscilla S S, Maria F A B, Ana P N P, Hugo C R J and Pericles B A 2011 *Industrial Crops and Products* **34** 831-837.

[6] Hasheimi S R, Zulkifli I, Somchit M N, Zunita Z, Loh T C, Soleimani A F and Tang S C 2013 *J. of Animal Physiology and Animal Nutrition* **97** 632–638.

[7] Ayssiwede, S. B., A. Dieng, H. Bello, C. A. A. M. Chrysostome, M.B. Hane, A. Mankor, M. Dahouda, M. R. Hounato, J. L. Hornick, A. Missohou. 2011. Effects of *Moringa oleifera* (*Lam.*) leaves meal incorporation in diets on growth performances, carcass characteristics and economics results of growing indigenous Senegal chickens. *Pak. J. Nutr.*, 10 (12): 1132-1145.

[8] Agustine A.A. and Ufuoma., 2013. Flavonoid from the leaves of *Physalis angulata L*, *Planta Medica.* 79 (13), 1211. alami sebagai pakan fungsional. *Wartazoa.* 23(1): 31-40.

[9] Dewi, T. K.,I. G.Bidura, dan D. P.Candrawati. 2014. The effect of giving moringa (*Moringa oleifera*) and garlic (*Allium sativum*) leaf extract through drinking water on performance of broiler age 2-6 weeks. *Peternakan Tropika.* Vol 2 (3): 461-475.

[10] Mboti, C. I., Eja, M. E., Adegoke, A. A., Iwaat, G. D., Asikong, B. E., Takon, I., Udo, S. M., and Akeh, M. 2009. Phytochemical Properties and Antimicrobial Activities of Combined effect of Ektracts of the Leaves of *Garcinia Kola*, *Vernonia*

**Commented [D7]:** Please consider Since there no statistical different, what kind of positive effects on broiler? Please describe it

amygdalina and Honey on Some Medically Important Microorganisms. *Sfr. J. Microbiol, Res.* 3(9):557-559.

- [11] Bharali, R. 2003. Chemomodulatory effect of moringa oleifera Lam. On hepatic carcinogen metabolizing enzymes, antioxidant parameters and skin papillomagenesis in mice. *Asian Pacific J Cancer Prevent* 4:2.
- [12] Amad , A.A., K. Manner, K. R. Wendler, K. Neumann and J. Zentek. 2011. Effects of a phytogenic feed additive on growth performance and ileal nutrient digestibility in broiler chickens. *Poultry Science.* 90 :2811–2816.
- [13] Yulrahmen, R. 2008. Khasiat dan Manfaat Minyak Atsiri Daun Sirih. Agromedia Puataka. Yogyakarta.
- [14] Dewi, T. K.,L. G,Bidura, dan D. P.Candrawati. 2014. The effect of giving moringa (*Moringa oleifera*) and garlic (*Allium sativum*) leaf extract through drinking water on performance of broiler age 2-6 weeks. *Peternakan Tropika.* Vol 2 (3): 461-475.
- [15] Daud M., Yaman M A, Usman Y and Y Aqmarina. 2021. Potential of ciplukan extract (*Physalis angulata* L.) and patchouli waste (pogostemon patchouli pellet) as alternative sources of phytogenic feed additive. *IOP Conf. Series: Earth and Environmental Science* 644 (2021) 012072.

- [16] Dzulknain B, D. Sundari dan A.Chozin. 1996. Tanaman obat bersifat antibakteri di Indonesia. *Cermin Dunia Kedokteran.* 110:35-48.