

Comparison of Exclusive and Non Exclusive Breast Feeding with the Incidence of Allergic Rhinitis in Children

by Benny Kurnia

Submission date: 09-Nov-2021 11:59AM (UTC+0700)

Submission ID: 1697478588

File name: Mulya_Safri,_et_al.pdf (458.68K)

Word count: 2186

Character count: 11923

doi: <https://doi.org/10.20546/ijcrar.2018.604.003>

6

Comparison of Exclusive and Non Exclusive Breast Feeding with the Incidence of Allergic Rhinitis in Children

Mulya Safri^{1*}, Muhammad Dipi Abdillah², Aulia Rahman Putra³, Benny Kurnia⁴ and Muhsin²

7

¹Departmen³⁵t of Child Health Syiah Kuala University, Dr. Zainoel Abidin Hospital, Banda Aceh, Indonesia

²Fac³⁷ulty of Medicine, Syiah Kuala University, Banda Aceh, Indonesia

³Dr. Zainoel Abidin Hospital, Banda⁷Aceh, Indonesia

⁴Department of Head Neck Surgery Syiah Kuala University, Dr. Zainoel Abidin Hospital, Banda Aceh, Indonesia

*Corresponding author

Abstract

Allergic rhinitis is common chronic disease in children and the incidence increases in children aged over³¹ years. Exclusive breastfeeding may reduce the risk of allergic rhinitis due to its anti-allergen and anti-inflammatory properties. This study aimed to find the difference between children who were exclusively breastfed and did not exclusively breastfeed²⁰ the incidence of allergic rhinitis in children. This analytic comparative with cross sectional study was conducted in Dr. Zainoel Abidin Hospital, Banda Aceh. Data collected from medical records and questionnaires completed by parents. A total of 35 children aged 5 to 18 years were collected using accidental sampling. The results shown 20 (57.1%) children had a history of exclusive breastfeeding and 15 (42.9%) children had a history of non-exclusive breastfeeding. About 22 (62.9 %) children suffering from intermittent allergic rhinitis, while other 13 (37.1 %) children suffering persistent allergic rhinitis. Fisher's exact²⁶ results shown statistically significant difference between Exclusive breastfeeding and non-exclusive breastfeeding to the incidence of allergic rhinitis in children aged 5 to 18 years ($P=0.005$; OR 6.9 95% CI 1.78-26.85). In conclusion, children with a history of exclusive breastfeeding had a lower frequency of allergic rhinitis symptoms. It is suggested²² that children with a moderate or high risk of allergy to be given exclusive breast feeding in the first 6 months of life to reduce the incidence of allergic rhinitis in the future.

Article Info

Accepted: 07 March 2017
Available Online: 20 April 2018

Keywords

Allergic rhinitis,
Exclusive breast feeding,
Children aged 5-18 years

Introduction

Allergic rhinitis is a chronic upper airway disease induced by allergen exposure and produces an inflammatory process mediated by IgE with symptoms of rhinorrhea, sneezing, nasal congestion, and/or itching of the nose or eye (Tharpe and Kemp, 2015; Mims 2012). Allergic Rhinitis and its Impact on Asthma (ARIA), in

collaboration with World Health Organization (WHO), states that allergic rhinitis represents a global health problem affecting around 10 to 20% of the population and this number continues year to year (Broek *et al.*, 2010). The results of the International Study of Asthma and Allergies in Childhood (ISAAC) surveys shown the prevalence of allergic rhinitis varies worldwide. Prevalence in Southeast Asia around 5.5% to 44.2%

25

(Tong and Lin, 2015). Allergic rhinitis is not common in children under 2 years of age, since exposure to allergens usually takes several years to develop (Tharpe and Kemp, 2015). Research conducted by ISAAC shown genetic (family history of allergy), age of child, gender, frequent airway infections, antibiotics in the first year of life, have pets, breastfeeding, socioeconomic status and the process of cesarean delivery are common risk factor for allergic rhinitis (Ji *et al.*, 2016; Li *et al.*, 2015).

Allergic rhinitis disease should be prevented as early as possible. Giving exclusive breastfeeding in the first 6 months of life should be a preventive factors against allergic rhinitis. (Wuryanti and Wibowo, 2015; Friedman *et al.*, 2005; Adkinson, 2008). Previous study shown that children who did not receive exclusive breastfeeding has a prevalence of allergic rhinitis 12.4% higher than children who consumed exclusive breastfeeding (Ehlayel *et al.*, 2008).

24

Materials and Methods

This analytic comparative with cross-sectional study was conducted in Dr. Zainoel Abidin Hospital Banda Aceh, the capital of Aceh Province, from September to December 2017. A total of 45 children aged 5 to 18 years were collected by accidental sampling. Children with a history of breastfeeding during infancy (aged 0 to 6 months) were included into inclusion criteria. Children suffering from infections and congenital problem were excluded in this study. Children had been diagnosed by an allergy and immunology specialist, based on examination and previous history of allergy and then divided into intermittent allergic rhinitis and persistent allergic rhinitis. Data collected from examinations, questionnaires completed by parents and medical records. Fisher's exact test were performed to analyze the obtained data. Data collected included age, sex, history of breastfeeding, and history of previous treatment. All analyses were performed with SPSS ver. 22.0 (IBM SPSS Statistics for Windows Ver 22.0. Armonk, NY, USA). Test results revealed statistically significant if *P* value of <0.05.

Results and Discussion

We obtained 45 children that fulfilled the inclusion criteria. Subject characteristics are shown in Table 1. Most of the subject aged between 5 to 11 years old (64.4%). In intermittent allergic rhinitis group, 81.2% of subjects feel slightly interrupted in performing daily activities, but in persistent allergic rhinitis group, 87.5%

of subjects feel very interrupted in performing daily activities. Exclusive breastfeeding during infancy (0 to 6 months) were given into 81.5% subjects in intermittent allergic rhinitis group. And in persistent allergic rhinitis group, 61.1% of subjects were given non exclusive breastfeeding during infancy. About 71.1% of subjects were grouped into low-moderate risk allergy.

Subject in intermittent allergic rhinitis group have slightly different numbers than in persistent allergic rhinitis group, 29 (64.4%) vs 16 (35.6%), respectively (Figure 1). 27 (60%) of total subjects were given exclusive breastfeeding during infancy (0 to 6 months) (Figure 2).

We analyzed the relation between variables in Table 2 using Fisher's exact test because chi-square test is not suitable. The result showed a statistically significant difference between allergic rhinitis and feeding history in infancy ($P=0.005$). We also conducted a Maentel-Haenszel analysis to find the risk of allergic rhinitis on a history of breastfeeding. We found that children who consumed non exclusive breastfeeding will have 6.9 times higher to suffer persistent allergic rhinitis (95% CI 1.78-26.85).

Allergic rhinitis disease should be prevented as early as possible because this problem will inhibit the growth and development process of children (Wuryanti and Wibowo, 2015). Giving exclusive breast feeding should be a preventive factors against allergic rhinitis. This type of milk contains Transforming growth factor (TGF- β), CD-14, lactoferrin, and IgA which plays an important role in the prevention of allergen exposure and the development of allergies in children. TGF- β improving the individual ability for producing inner IgA against substances that can trigger allergy as β -lactoglobulin, casein, gliadin, and ovalbumin that are normally on food or drinks. TGF- β also inhibits the inflammatory process of T helper-2 cells, modulates antigens presenting cell (APC) and induces IgA production. High level of CD-14 content in breast milk will play a protective function from allergic development. Lactoferrin helps to accelerate gastrointestinal maturation that can prevent allergies. Secretory IgA in breast milk also helps the intestine from allergic events by coating the intestinal barrier mucosa to prevent the occurrence of allergen sensitization (Friedman *et al.*, 2005; Adkinson, 2008). Ehlayel *et al.*, (2008) states that children who did not receive exclusive breastfeeding has a prevalence of allergic rhinitis 12.4% higher than children who consumed exclusive breastfeeding.

Table.1 Subject characteristics

Variable	Intermittent allergic rhinitis		Persistent allergic rhinitis		Total n (%)
	n	%	n	%	
Gender					
Male	15	65.2	8	34.8	23 (51.1)
Female	14	63.6	8	36.4	22 (48.9)
Age					
Child (5 to 11 years)	20	69	9	31	29 (64.4)
Early adolescent (12 to 16 years)	8	57.1	6	42.9	14 (31.2)
Teenager (17 to 18 years)	1	50	1	50	2 (4.4)
Interrupted activity					
Not interrupted	2	40	3	60	5 (11.1)
Slightly interrupted	26	81.2	6	18.3	32 (71.1)
Very interrupted	1	12.5	7	87.5	8 (17.8)
Allergy Risk					
Low-moderate	23	71.9	9	28.1	32 (71.1)
High	6	46.2	7	53.8	13 (28.9)

Table.2 Bivariate analyzed between allergic rhinitis and breast feeding

Variable	Allergic rhinitis		P	OR	95% CI
	Intermittent: n (%)	Persistent: n (%)			
Feeding history					
Exclusive breastfeeding	22 (81.5)	5 (18.5)	0.005	6.9	1.78-26.85
Non exclusive breastfeeding	7 (38.9)	11 (61.1)			



Fig.1 Comparison of intermittent and persistent allergic rhinitis

Fig.2 Comparison of exclusive and non exclusive breast feeding



Similar in our results, allergic rhinitis and feeding history in infancy shown a statistically significant difference where children who consumed non exclusive breast feeding will have 6.9 times higher to suffer persistent allergic rhinitis ($P=0.005$; 95% CI 1.78-26.85).

Allergic rhinitis does not cause the death but can affect the quality of life because of sleep disturbance, decreased concentration and work productivity, decreased performance in school and social activities and even psychological disorders such as depression if allergicsymptom becomeworsening (Rotiroti and Scadding, 2016; Wuryanti and Wibowo, 2015).

Allergic rhinitis in fact disturbs the patient in performing daily activities. About 81% of subjects suffering from intermittent allergic rhinitis experienced slightly interrupted in performing daily activities, while 87% of patients who had persistent allergic rhinitis experienced severe interrupted inperforming daily activities.

This is in accordance with study conducted by Eli O. Meltzer (2016) which states that allergic rhinitis can affect the quality of life in a person including physical and socialdaily activities, sleep disturbances, fatigue, irritability, depression, attention deficit, memory and learningdisturbance.

Allergy symptoms are also related to genetics. This symptoms were found in 39.8% of children if their mother has a history of allergy and 30.2% if their father has the symptoms of allergy (Johansson *et al.*, 2004) ³³ e divide allergic risk factors into low-moderate and high risk of allergy based on family history. 72% of allergic rhinitis children were included into low-moderate group.

In conclusion, Allergic rhinitis does not cause the death but can affect the quality of life especially make the sufferer experienced sleep disturbances, fatigue, irritability, depression, attention deficit, memory and learning disturbances. Allergic rhinitis and feeding history in infancy showed a relation where children who consumed non exclusive breastfeeding will have 6.9 times higher to suffer persistent allergic rhinitis. It is highly recommended to be given exclusive breastfeeding during infancyasa preventive factors against allergic rhinitis.

References

- ¹⁶ Adkinson NF. Middleton's Allergy: Principles and Practice 7th Edition. 7th ed. Philadelphia: Mosby Elsevier; 2008.
- ²⁸ Broek JL, Bousquet J, Baena-Cagnani CE, Bonini S, Canonica GW, Casale TB, *et al.*, Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines: 2010 Revision. J Allergy Clin Immunol. 2010;126(3): 466–76.
- ⁵ Ehlayel MS, Bener A. Duration of breast-feeding and the risk of childhood allergic disease in a developing country. Allergy Asthma Proceed 2008; 29: 386-91.
- ³ Friedman NJ, Zeiger RS, Diego S. The Role of Breast-feeding in the Development of Allergies and Asthma. J Allergy Clin Immunol 2005;115(6): 1238–48.
- ¹¹ Ji Y, Liu Y, Yang N. Pediatric Rhinitis Risk Factors. Exp Ther Med 2016;12(4):2383–6.
- ⁸ Johansson SG, Bieber T, Dahl R, Friedmann PS, Lanier BQ, Lockey RC, *et al.*, Revised nomenclature for allergy for global use: report of the nomenclature review committee of the world allergy organization. J

- 17 Allergy Clin Immunol 2004;11:38-83.
- Li Y, Jiang Y, Li S, Shen X, Liu J, Jiang F. Pre-and Postnatal Risk Factors in Relation to Allergic Rhinitis in School-Aged Children in China. PLoS ONE 2015; 10(2): 1–11.
- 9 Meltzer EO. Allergic Rhinitis Burden of illness, Qualify of life Comorbidities and Control. Immunol Allergy Clin NAm 2016; 36: 235-48.
- 14 Mims JW. Allergic Rhinitis. Facial Plast Surg Clin North Am 2012; 20(1): 11-20.
- 10 Rotiroti G, Scadding GK. Allergic Rhinitis-an Overview of a Common Disease. Paediatr Child Health 2016; 26(7): 298–303.
- 12 Tharpe CA, Kemp SF. Pediatric Allergy Rhinitis. Immunol Allergy Clin North Am 2015; 35(1):185-98.
- 29 Tong M, Lin J. Epidemiology of Allergic Rhinitis throughtout the world. In: Akdis CA, Hellings PW, Agache I, editors. Global Atlas of Allergic Rhinitis and Chronic Rhinosinusitis by European Acedemy of Allergy and Clinical Immunology. European Academy of Allergy and Clinical Immunology 2015; p. 62.
- 15 Wuryanti T, Wibowo T. ASI Eksklusif sebagai Faktor Protektif Rinitis Alergi pada Anak. Sari Pediatr 2015;17(1):59–63.

How to cite this article:

Mulya Safri, Muhammad Dipi Abdillah, Aulia Rahman Putra, Benny Kurnia and Muhsin. 2018. Comparison of Exclusive and Non Exclusive Breast Feeding with the Incidence of Allergic Rhinitis in Children. *Int.J.Curr.Res.Aca.Rev.* 6(4), 15-19. doi: <https://doi.org/10.20546/ijcrar.2018.604.003>

Comparison of Exclusive and Non Exclusive Breast Feeding with the Incidence of Allergic Rhinitis in Children

ORIGINALITY REPORT

24%
SIMILARITY INDEX

20%
INTERNET SOURCES

17%
PUBLICATIONS

9%
STUDENT PAPERS

PRIMARY SOURCES

1 In collaboration with the World Health Organization. "Allergic Rhinitis and Its Impact on Asthma", The Journal of Allergy and Clinical Immunology, 200111 1%
Publication

2 fsd.unsyiah.ac.id 1%
Internet Source

3 scholarworks.utrgv.edu 1%
Internet Source

4 mafiadoc.com 1%
Internet Source

5 pubmed.ncbi.nlm.nih.gov 1%
Internet Source

6 sinta3.ristekdikti.go.id 1%
Internet Source

7 www.balimedicaljournal.org 1%
Internet Source

8 www.webmedcentral.com
Internet Source

1 %

9

K. Trikojat, A. Buske-Kirschbaum, F. Plessow, J. Schmitt, R. Fischer. "Memory and multitasking performance during acute allergic inflammation in seasonal allergic rhinitis", Clinical & Experimental Allergy, 2017

Publication

1 %

10

Yvette N. Lamb. "Olopatadine/mometasone combination nasal spray in allergic rhinitis: a profile of its use", Drugs & Therapy Perspectives, 2020

Publication

1 %

11

www.verywellhealth.com

Internet Source

1 %

12

Logan Christensen, John T. McDonnell, Jasleen Singh. "Chapter 3 Ocular Manifestations of Allergic and Immunologic Diseases", Springer Science and Business Media LLC, 2017

Publication

1 %

13

G Berkiten, İ Aydoğdu, T L Kumral, Z Saltürk, Y Uyar, A Arslanoğlu, B Tutar, A E Göker. "Nasal eosinophilia in nasal smears of patients with persistent and intermittent allergic rhinitis", The Journal of Laryngology & Otology, 2018

Publication

1 %

14

www.wakehealth.edu

Internet Source

1 %

15

doaj.org

Internet Source

1 %

16

link.springer.com

Internet Source

1 %

17

www.wjgnet.com

Internet Source

1 %

18

journals.ashs.org

Internet Source

1 %

19

Submitted to University of Melbourne

Student Paper

<1 %

20

eprints.uanl.mx

Internet Source

<1 %

21

query.library.utoronto.ca

Internet Source

<1 %

22

www.dovepress.com

Internet Source

<1 %

23

www.fantaproject.org

Internet Source

<1 %

24

www.medigraphic.com

Internet Source

<1 %

25

J. Bousquet. "Executive Summary of the Workshop Report 7-10 December 1999,

<1 %

26	etd.repository.ugm.ac.id	<1 %
	Internet Source	

27	test.dovepress.com	<1 %
	Internet Source	

28	www.researchsquare.com	<1 %
	Internet Source	

29	Submitted to Erasmus University Rotterdam	<1 %
	Student Paper	

30	Kezhi Liu, Xuemei Liang, Lanting Guo, Yuan Li, Xirong Li, Bo Xin, Mingjin Huang, Yuanyuan Li. "The acute stress disorder in the paediatric surgical children and adolescents injured in the Wenchuan earthquake of China", Stress and Health, 2010	<1 %
	Publication	

31	Na Liu, Jitu Wang, Sainan Qiu, Man Zhang. "Changes of Serum Bilirubin and Urine Peptides Related to Bilirubin Metabolism in Patients with Allergic Rhinitis", Research Square Platform LLC, 2021	<1 %
	Publication	

32	Youjin Li, Yanrui Jiang, Shenghui Li, Xiaoming Shen, Jinfen Liu, Fan Jiang. "Pre- and Postnatal Risk Factors in Relation to Allergic Rhinitis in	<1 %
----	--	------

School-Aged Children in China", PLOS ONE, 2015

Publication

33

ctajournal.biomedcentral.com

Internet Source

<1 %

34

www.ijss-sn.com

Internet Source

<1 %

35

www.paom.pl

Internet Source

<1 %

36

J. Bousquet. "Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 Update (in collaboration with the World Health Organization, GA²LEN and AllerGen**)", *Allergy*, 4/2008

Publication

<1 %

37

Cut Husna. "Chapter 2 Acute Respiratory Care and Wound Care of Tsunami Patients: What Should the Nurses Do?", IntechOpen, 2012

Publication

<1 %

38

Tamay, Z.. "Prevalence and risk factors for allergic rhinitis in primary school children", *International Journal of Pediatric Otorhinolaryngology*, 200703

Publication

<1 %

Exclude quotes On

Exclude bibliography Off

Exclude matches Off